



## Maternal and Child Health in the Context of Nutrition, Oral Health, and Emergency Care: An Integrated Nursing, Dentistry, Social Work, Public Health, Epidemiology, and Emergency Medicine Perspective

Salha Mohsen Hassan <sup>(1)</sup>, Mihammed Hussien Dammas Matkooor <sup>(2)</sup>, Reem Abdu Ibrahim Gharawi <sup>(3)</sup>, Fawaz Abdullah Alshammari <sup>(4)</sup>, Naser Saad Rashid Al Dosari <sup>(5)</sup>, Sultan Mashan Matlaq Alotaibi <sup>(6)</sup>, Abdullah Awad Butihan Alotaibi <sup>(6)</sup>, Ayat Abdullah Alghwinem <sup>(7)</sup>, Abdulrahman Samir Khateiri <sup>(8)</sup>, Fahad Hamad Hassan Alqahtani <sup>(9)</sup>, Fatimah Mohammed Hassan Duryib <sup>(10)</sup>, BeSheah Ali Samaan Zubaidi <sup>(10)</sup>, Adel Farhan Alanazi <sup>(11)</sup>, Bandar Abdullah Khalifah Alanazi <sup>(12)</sup>

(1) King Fahd Central Hospital, Ministry of Health, Saudi Arabia,

(2) Al-Mawsim General Hospital, Ministry of Health, Saudi Arabia,

(3) Al-Mawsim General Hospital, Jazan Health Cluster, Ministry of Health, Saudi Arabia,

(4) Al-Uwayqilah General Hospital, Ministry of Health, Saudi Arabia,

(5) Wadi Al Dawasir General Hospital, Ministry of Health, Saudi Arabia,

(6) Sajer General Hospital, Ministry of Health, Saudi Arabia,

(7) Prince Saud Bin Jalawi Hospital, Ministry of Health, Saudi Arabia,

(8) King Fahad Hospital, Madinah, Ministry of Health, Saudi Arabia,

(9) Dhahran Eye Specialist Hospital, Ministry of Health, Saudi Arabia,

(10) Al-Malha Primary Health Care Center, Western Sector, Jazan, Ministry of Health, Saudi Arabia,

(11) Al-Muzahmiya Hospital, Riyadh, Ministry of Health, First Health Cluster, Saudi Arabia,

(12) Ministry of Health Office in Alqurayyat, Saudi Arabia

### Abstract

**Background:** Maternal and child health (MCH) disparities persist globally, driven by interconnected factors including poor nutrition, dental neglect, and inadequate emergency preparedness. Traditional siloed care delivery fails to address these intersecting determinants. **Aim:** This narrative review synthesizes evidence on integrated MCH interventions spanning prenatal/postnatal nursing, maternal oral health and early childhood caries prevention, family support for food insecurity, public health nutrition programs, epidemiological surveillance of malnutrition, and obstetric/pediatric emergency care. **Methods:** A comprehensive literature search (2015-2024) was conducted across nursing, dental, public health, and emergency medicine databases. **Results:** Integrated care models reduce MCH disparities through: nurse-led home visiting programs (30-50% reduction in adverse outcomes); dental-nursing partnerships (40% early childhood caries reduction); social work-nutrition collaborations (25-35% food insecurity mitigation); and emergency medicine-epidemiology linkages (improved obstetric triage and pediatric trauma response). **Conclusion:** Interprofessional MCH integration addressing nutrition, oral health, and emergency care effectively reduces disparities. Policy recommendations include reimbursement reform and cross-sector workforce development.

**Keywords:** maternal-child health; integrated care; nutrition; oral health; emergency care

### Introduction

Maternal and child health (MCH) represents a fundamental indicator of population health and healthcare system effectiveness, yet persistent disparities across socioeconomic, racial, and geographic lines demonstrate that conventional siloed care delivery models are insufficient to address the complex, multifactorial determinants of MCH outcomes (Braveman & Gottlieb, 2014). The health of mothers and children does not exist in isolation but rather emerges from the dynamic interaction of biological, behavioral, social, and healthcare system factors that span multiple professional domains. Nutrition status during pregnancy influences both

maternal outcomes (gestational diabetes, hypertensive disorders) and child development (low birth weight, neurocognitive deficits). Oral health, often neglected in prenatal care, has established associations with adverse pregnancy outcomes including preterm birth and low birth weight, while early childhood caries (ECC) remains the most common chronic disease of childhood, disproportionately affecting low-income and minority populations (Wade et al., 2019).

Emergency care for obstetric and pediatric populations represents the safety net for failures in preventive and primary care. Obstetric emergencies, including hemorrhage, eclampsia, and sepsis, account for substantial maternal morbidity and mortality,

while pediatric trauma and acute illness frequently result from preventable conditions linked to poverty, malnutrition, and inadequate supervision (American Academy of Pediatrics, 2018). The COVID-19 pandemic exposed and exacerbated these vulnerabilities, disrupting nutrition programs, dental services, and routine preventive care while increasing food insecurity and delaying emergency care seeking (Wolfson et al., 2021).

The recognition that these challenges are inherently cross-disciplinary has driven the development of integrated care models that bring together nursing (prenatal and postnatal home visiting), dentistry (maternal oral health and ECC prevention), social work (family support and food insecurity case management), public health (nutrition programs including WIC and SNAP), epidemiology and health inspection (surveillance of malnutrition and outbreak detection), and emergency medicine (obstetric emergencies and pediatric trauma response) (Brashers et al., 2015). This narrative review synthesizes current evidence on integrated MCH interventions addressing nutrition, oral health, and emergency care, with the explicit aim of identifying effective strategies for reducing disparities through interprofessional collaboration. The central thesis is that integrated, team-based care spanning the prenatal-to-early childhood continuum, delivered through coordinated nursing-dentistry-social work-public health-epidemiology-emergency medicine partnerships, can substantially reduce MCH disparities compared to traditional siloed approaches.

### Methodology

This narrative review followed established guidelines for synthesizing evidence across multiple disciplines. A comprehensive literature search was conducted across PubMed/MEDLINE, CINAHL (nursing), Dentistry & Oral Sciences Source, Social Work Abstracts, Public Health Database, and Web of Science for studies published between January 2015 and December 2024. Search strategies combined terms for maternal-child health, nutrition, oral health, emergency care, nursing, dentistry, social work, public health, and epidemiology with interprofessional collaboration and integrated care concepts. Inclusion criteria were: (1) peer-reviewed original research, systematic reviews, or evidence-based guidelines; (2) focus on MCH outcomes related to nutrition, oral health, or emergency care; (3) explicit attention to interprofessional or integrated care approaches; (4) published in English; (5) from 2015-2024. Exclusion criteria were single-discipline studies without integration components, non-human studies, and opinion pieces without empirical data. Data extraction focused on intervention characteristics, integration mechanisms, outcomes, and disparity reduction effects. Given the narrative design, findings were synthesized thematically across professional domains.

### Prenatal and Postnatal Care Integration

### The Nurse Home Visiting Model as an Integration Platform

Nurse home visiting programs represent the most extensively evaluated platform for integrated MCH care delivery. The Nurse-Family Partnership (NFP), which provides structured nurse home visits from pregnancy through the child's second birthday, has demonstrated consistent effects on MCH outcomes across multiple randomized controlled trials (Odeny et al., 2014). The mechanism of effect extends beyond traditional nursing care to include coordination across nutrition services, dental referrals, social work support, and emergency care navigation. Nurses in integrated models receive training to screen for food insecurity, conduct oral health assessments using standardized tools, identify warning signs of obstetric emergencies, and facilitate warm handoffs to interdisciplinary team members.

A recent systematic review of 23 nurse home visiting programs found that integrated models achieved a 32% reduction in preterm birth among high-risk mothers and a 28% reduction in emergency department visits for non-emergent conditions (Molloy et al., 2021). The integration of standardized screening protocols for social determinants of health—including the Hunger Vital Sign for food insecurity and the Oral Health Basic Screening Survey for pregnant women—enabled nurses to identify needs that would otherwise remain unrecognized until crisis presentation. Programs that embedded social workers within nurse home visiting teams achieved significantly higher rates of food assistance enrollment (74% vs. 41% in referral-only models) (Jack et al., 2019).

### Postnatal Nursing Support for Breastfeeding and Infant Nutrition

Postnatal nursing care provides critical opportunities for nutrition intervention that reduce disparities in breastfeeding initiation and duration. Breastfeeding confers protective effects against ECC, obesity, and acute infectious diseases, yet rates are substantially lower among low-income and minority populations (Meek & Noble, 2022). Integrated nursing programs that include lactation consultants, dental hygienists, and nutritionists achieve breastfeeding duration rates 40% higher than standard care at 6 months postpartum. The integration of oral health education into postnatal breastfeeding support—including guidance on oral hygiene following feeds and early introduction of oral care practices—has been associated with delayed colonization by cariogenic bacteria and reduced ECC risk at 12 months (Edelstein, 2017).

Nurse-led postpartum depression screening, when integrated with social work follow-up, addresses the bidirectional relationship between maternal mental health and child nutrition. Depressed mothers are 2.5 times more likely to report food insecurity and 3 times more likely to discontinue breastfeeding before 6 months (Slomian et al., 2019). Integrated care models

that co-locate mental health screening, social work case management, and nutrition counseling within postnatal nursing visits achieve 55% higher resolution of maternal depression and 42% higher sustained breastfeeding rates compared to fragmented referral systems (Tandon et al., 2020).

**Maternal Oral Health and Early Childhood Caries The Pregnancy as a Teachable Moment**

Pregnancy represents a "teachable moment" for oral health behavior change, yet dental care during pregnancy remains underutilized, particularly among Medicaid-enrolled and uninsured women (Vamos et al., 2023). Integrated nursing-dental models have demonstrated success in overcoming barriers, including provider reluctance, patient misconceptions about dental treatment safety during pregnancy, and lack of referral systems (Table 1). The integration of oral health risk assessment into routine prenatal nursing visits—using tools such as the Oral Health Risk Assessment Tool (OHRAT) and the Pregnancy Risk Assessment Monitoring System (PRAMS) oral health module—identifies women requiring dental referral and provides brief motivational interviewing to address perceived barriers.

A cluster-randomized trial of an integrated prenatal oral health program involving nurse-delivered oral health education, dental hygienist co-visits, and social work-facilitated transportation and childcare assistance found a 47% increase in dental attendance during pregnancy and a 38% reduction in ECC at 24 months (Wade et al., 2019). Women receiving integrated care were 3.2 times more likely to receive needed dental treatment during pregnancy compared to women receiving usual care (referral without navigation support). The program was most

effective for Medicaid-enrolled women, reducing the disparity in dental attendance between privately insured and Medicaid-enrolled women from 41% to 12%.

**Early Childhood Caries Prevention through Interprofessional Collaboration**

ECC is preventable yet remains the most common chronic disease of early childhood, with a prevalence of 23% among children aged 2-5 years and rates exceeding 40% among low-income and minority populations (Tinanoff et al., 2019). The integration of dental prevention into well-child visits, nurse home visits, and WIC nutrition appointments represents an evidence-based strategy for reducing ECC disparities. The "First Tooth" model, implemented in multiple states, trains non-dental professionals (nurses, WIC nutritionists, social workers) to conduct oral health risk assessments, apply fluoride varnish, provide anticipatory guidance, and facilitate dental referral (Braun et al., 2024).

Evaluation of integrated models demonstrates that nurse-administered fluoride varnish programs achieve 34-45% reductions in ECC incidence when implemented as part of routine well-child care (Linden et al., 2023). Social work integration addressing family-level barriers—including parental oral health literacy, competing priorities, and access to dental insurance—increases completion of dental referrals from 28% in standard care to 71% in integrated models (Edelstein, 2017). Public health dental hygienists practicing in collaborative agreements with nurses and social workers have extended ECC prevention to populations with limited dental access, including rural and migrant farmworker communities (Adeghe, 2024).

**Table 1: Integrated Nursing-Dental Interventions for Maternal and Child Oral Health Disparities Reduction**

Intervention Model	Professional Integration	Target Population	Key Outcome	Disparity Reduction
Nurse-delivered oral health assessment + dental navigator	Nursing, Dentistry, Social Work	Medicaid-enrolled pregnant women	47% increase in dental attendance; 38% reduction in ECC at 24 months	Reduced disparity in dental attendance: 41% to 12% (Wade et al., 2019)
Fluoride varnish in WIC settings	Public Health Nutrition, Dental Hygiene	Low-income children 6-36 months	41% reduction in ECC incidence	Reduced disparity between WIC and non-WIC children: 28% to 9% (Linden et al., 2023)
Nurse home visiting + dental referral coordination	Nursing, Social Work, Dentistry	First-time, low-income mothers	71% dental referral completion vs. 28% standard care	Eliminated racial disparity in dental referral completion (Edelstein, 2017)
Co-located dental hygienist in pediatric primary care	Dentistry, Nursing, Public Health	Children 0-5 years, all payers	82% of children receive preventive dental care by age 1 (vs. 22% nationally)	Reduced SES gradient in dental care receipt (Adeghe, 2024)
Tele-dentistry + nurse case management	Dentistry, Nursing, Social Work	Rural and frontier populations	56% reduction in emergent dental visits; 44% reduction in ED	Eliminated geographic disparity in preventive dental access (Braun et al., 2024)

---

use for dental  
conditions

---

### **Family Support and Food Insecurity Food Insecurity Screening and Intervention**

Food insecurity—limited or uncertain access to adequate food—affects 13.8 million US households with children and is associated with adverse maternal and child health outcomes including gestational diabetes, low birth weight, developmental delay, and increased emergency department use (Gundersen & Ziliak, 2018). Integrated social work-nutrition programs have demonstrated effectiveness in identifying food-insecure families and connecting them to federal nutrition assistance programs including the Supplemental Nutrition Assistance Program (SNAP), the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC), and school meal programs.

Universal food insecurity screening in prenatal clinics, pediatric primary care, and emergency departments, when combined with social work-facilitated enrolment assistance, increases SNAP and WIC participation by 50-70% compared to referral-only approaches (De Marchis et al., 2019). Social work integration within nurse home visiting programs achieves 85% enrolment of eligible families in WIC, compared to 45% in standard care (Jack et al., 2019). The mechanism of effect includes addressing documentation barriers, transportation to enrolment sites, and competing priorities—factors that disproportionately affect low-income and immigrant families. Longitudinal studies demonstrate that integrated food insecurity interventions reduce maternal depression (31% reduction), improve child dietary quality (28% improvement in Healthy Eating Index scores), and decrease emergency department visits for nutrition-sensitive conditions, including failure to thrive and dehydration (Gebremeskel et al., 2022).

### **The Intersection of Food Insecurity, Oral Health, and Emergency Care**

Food insecurity and poor oral health are bidirectionally related and synergistically increase emergency care utilization. Children with both food insecurity and untreated dental caries have 4.2 times higher odds of emergency department visits for dental conditions and 2.8 times higher odds of hospitalization for nutrition-sensitive conditions compared to food-secure children without caries (Bahanan et al., 2021). Social work-led integrated case management addressing both food insecurity and dental access reduces this dual burden, achieving 55% reduction in combined food insecurity-dental emergency visits (Thomas et al., 2019).

The integration of social workers into emergency departments for MCH populations—a model known as emergency department-based case management—has demonstrated effectiveness in

reducing recurrent visits for preventable conditions. Social workers in this role conduct food insecurity screening, facilitate SNAP/WIC enrollment, arrange dental appointments, and coordinate follow-up with primary care, nursing home visiting, and public health nutrition programs (Balasa et al., 2023). A randomized controlled trial of emergency department-integrated social work for families with children aged 0-3 years found a 37% reduction in 90-day emergency department revisits, 42% reduction in unmet food needs, and 34% reduction in unmet dental needs compared to standard emergency department care (Burnett et al., 2023).

### **Nutrition Programs and Population-Level Interventions**

#### **WIC and the Integration of Nutrition, Oral Health, and Nursing**

The Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) serves approximately 6.2 million participants monthly, providing nutrition education, supplemental foods, breastfeeding support, and healthcare referrals (Ventura et al., 2022). WIC represents an underutilized platform for integrated MCH care, given its reach to low-income pregnant and postpartum women, infants, and children up to age 5. Integration of oral health screening and fluoride varnish into WIC appointments has been implemented in multiple states, with evaluation demonstrating 41% reduction in ECC among participating children (Linden et al., 2023). Integration of nurse home visiting referrals from WIC increases enrollment in evidence-based home visiting programs by 35% compared to WIC alone (Odeny et al., 2014).

WIC-based breastfeeding peer counseling programs, when integrated with dental hygienist-delivered oral health education for mothers, achieve breastfeeding duration rates at 6 months of 58% compared to 42% in standard WIC (Meek & Noble, 2022). The integration of social work services within WIC—including assistance with housing, utilities, and transportation—addresses competing priorities that limit participation and adherence to nutrition recommendations. WIC sites with co-located social workers achieve 89% retention rates compared to 67% in standard WIC sites, with corresponding improvements in child immunization rates (15% increase) and lead screening completion (22% increase) (De Marchis et al., 2019).

#### **School Meal Programs and Nutrition Surveillance**

The National School Lunch Program (NSLP) and School Breakfast Program (SBP) serve over 30 million children daily, representing critical safety net nutrition programs that reduce food insecurity and improve dietary quality (Gundersen & Ziliak, 2018). Integrated public health-epidemiology-school

partnerships have enhanced the nutritional quality of school meals through menu reformulation, Farm to School programs, and nutrition education. Universal free school meal policies, implemented in several states and districts, reduce food insecurity by 34% and improve school attendance by 12% (Ruffini, 2022).

Epidemiological surveillance of child malnutrition, including monitoring of body mass index (BMI) trends and identification of emerging nutrition problems (e.g., micronutrient deficiencies, food insecurity hotspots), requires integration across public health, school health, and primary care. The integration of school nurse BMI screening with public health nutrition surveillance systems enables targeted interventions for children with obesity (referral to family-based weight management programs) and underweight (referral to WIC or SNAP and evaluation for underlying medical conditions) (Bejster et al., 2020). Health inspector integration ensures that school meal programs comply with nutritional standards and food safety requirements, preventing foodborne outbreaks that disproportionately affect young children (Paez & Alcorn, 2019).

#### **Surveillance of Malnutrition and Outbreaks Malnutrition Surveillance Systems**

Epidemiological surveillance of maternal and child malnutrition requires integration across healthcare, public health, and social service data systems. The Pregnancy Risk Assessment Monitoring System (PRAMS) and the Pediatric Nutrition Surveillance System (PedNSS) provide state-level data on nutrition-related indicators, including gestational weight gain, breastfeeding, food insecurity, and childhood obesity (Paez & Alcorn, 2019). Integration of PRAMS data with vital statistics, hospital discharge data, and emergency department visit data enables identification of populations at highest risk for adverse nutrition-related outcomes and evaluation of intervention effectiveness.

Health inspectors—professionals who monitor food safety in restaurants, schools, childcare facilities, and WIC clinics—play an underrecognized role in MCH nutrition protection. Integrated inspection-epidemiology programs have reduced foodborne illness outbreaks in childcare settings by 45% through targeted inspection protocols for high-risk facilities and rapid outbreak response protocols that include pediatric emergency department notification (Paez & Alcorn, 2019). The integration of health inspectors with WIC and school meal program nutritionists ensures that nutrition assistance is delivered through safe food systems, preventing the paradoxical outcome of foodborne illness from safety net programs (White et al., 2023).

#### **Outbreak Detection and Emergency Response**

Epidemiologic surveillance for outbreaks of infectious diseases—including vaccine-preventable diseases (measles, pertussis), gastrointestinal pathogens (norovirus, rotavirus), and emerging threats (COVID-19, respiratory syncytial virus)—requires

integration with emergency medicine for timely response. Maternal and child populations are disproportionately affected by outbreaks due to immature or compromised immune systems (infants, pregnant women) and congregate living settings (childcare, schools) (Gharpure et al., 2021). Integrated epidemiology-emergency medicine-outbreak response systems have demonstrated effectiveness in reducing outbreak-associated morbidity and mortality through early detection (epidemiology), rapid treatment (emergency medicine), and preventive interventions (public health).

The integration of emergency department syndromic surveillance with public health epidemiology systems enables the detection of outbreak signals days to weeks before traditional laboratory-based reporting. During the 2022 pediatric hepatitis of unknown etiology outbreak, integrated surveillance systems identified cases 10 days earlier than standard reporting, enabling rapid investigation and communication (Patel et al., 2022). For MCH populations, integrated epidemiology-emergency medicine systems have improved outcomes for maternal sepsis (24% reduction in mortality), pediatric severe malnutrition (35% reduction in hospitalization), and vaccine-preventable disease outbreaks (reduction in outbreak duration by 40%) (Paez & Alcorn, 2019).

#### **Obstetric Emergencies and Pediatric Trauma**

##### **Obstetric Emergency Preparedness and Response**

Obstetric emergencies—including hemorrhage, hypertensive emergencies (eclampsia), sepsis, thromboembolism, and cardiac events—account for a substantial proportion of severe maternal morbidity and mortality, with significant disparities by race, ethnicity, and socioeconomic status (American Academy of Pediatrics, 2018). Integrated emergency medicine-obstetric-nursing protocols, including obstetric emergency drills, hemorrhage carts, and massive transfusion protocols, have reduced maternal mortality from obstetric hemorrhage by 35% in hospitals implementing comprehensive preparedness programs (Imansari et al., 2023).

The integration of emergency medical services (EMS) with community health workers and nurse home visiting programs has improved prehospital care for obstetric emergencies (Table 2). Community paramedicine programs that train paramedics in obstetric emergency assessment and treatment, combined with community health worker-facilitated prenatal engagement, reduce the proportion of out-of-hospital births from 1.4% to 0.6% and reduce maternal transport time for hemorrhage by 22 minutes (Hill et al., 2023). The integration of social work into obstetric emergency care—including assistance with insurance, transportation, and childcare—addresses barriers to follow-up care that contribute to disparities in maternal outcomes (De Marchis et al., 2019).

##### **Pediatric Trauma and Acute Illness**

Pediatric trauma is the leading cause of death among children aged 1-19 years, with disparities in injury rates (3-fold higher for low-income children), prehospital care, and trauma center access (Sanchez et al., 2020). Integrated emergency medicine-public health-nursing injury prevention programs, including car seat distribution and education, home safety visits, and drowning prevention, have reduced pediatric injury mortality by 28% in implementing communities (Zonfrillo et al., 2014). The integration of social work into pediatric emergency care for injury-related visits enables identification of underlying social determinants (unsafe housing, lack of car seats, food insecurity, child maltreatment risk) and connection to preventive services.

Acute pediatric illness—including respiratory infections (bronchiolitis, pneumonia),

gastroenteritis with dehydration, and asthma exacerbations—accounts for over 20 million emergency department visits annually (Portillo et al., 2023). Integrated emergency medicine-nursing-public health programs have reduced avoidable emergency department visits through: (1) nurse telephone triage lines providing home management guidance; (2) social work-facilitated connection to primary care and nutrition programs; and (3) public health surveillance to identify and address community-level drivers (e.g., lead poisoning, air quality, food insecurity). The integration of dental emergencies into pediatric emergency care—including antibiotic and pain management with social work-facilitated dental follow-up—reduces repeat emergency department visits for dental conditions by 56% (Braun et al., 2024).

**Table 2: Integrated Emergency Medicine-MCH Interventions for Disparity Reduction**

Emergency Condition	Integrated Intervention	Professional Integration	Outcome	Disparity Impact
<b>Obstetric hemorrhage</b>	Obstetric emergency drills + hemorrhage cart + massive transfusion protocol	Emergency Medicine, Obstetrics, Nursing	35% reduction in maternal mortality	Reduced racial disparity in hemorrhage mortality from 3:1 to 1.5:1 (Imansari et al., 2023)
<b>Obstetric transport</b>	Community paramedicine + community health worker prenatal engagement	EMS, Health, Work	22-minute reduction in transport time; 57% reduction in out-of-hospital births	Reduced rural-urban disparity in maternal transport time (Hill et al., 2023)
<b>Pediatric trauma</b>	Car seat distribution + home safety visits + injury prevention education	Emergency Medicine, Public Health, Nursing	28% reduction in pediatric injury mortality	Reduced SES disparity in injury mortality from 3:1 to 1.8:1 (Zonfrillo et al., 2014)
<b>Pediatric dental emergency</b>	Pain management + antibiotics + social work-facilitated dental follow-up	Emergency Medicine, Dentistry, Social Work	56% reduction in repeat ED visits for dental conditions	Eliminated racial disparity in dental ED revisits (Braun et al., 2024)
<b>Child food insecurity</b>	ED-based food insecurity screening + social work-facilitated SNAP/WIC enrollment	Emergency Medicine, Social Work, Public Health	42% reduction in unmet food needs; 37% reduction in ED revisits	Reduced disparity in food insecurity between Medicaid and privately insured (Burnett et al., 2023)

### **Integrated Care Models Reducing Disparities: Synthesis and Mechanisms**

#### **The Interprofessional Collaborative Practice Framework**

The evidence synthesized across nursing, dentistry, social work, public health, epidemiology, and emergency medicine demonstrates that integrated care models consistently reduce MCH disparities compared to traditional siloed approaches. The Interprofessional Collaborative Practice (IPCP) framework—which includes core competencies in values/ethics, roles/responsibilities, interprofessional communication, and teams/teamwork—provides an organizing structure for understanding effective integration (Brashers et al., 2015). Successful

integrated MCH programs share common elements: (1) co-location of services or formal referral agreements with warm handoffs; (2) shared electronic health records or data sharing agreements; (3) cross-training of professionals in each other's core functions; (4) regular interprofessional team meetings; and (5) shared quality improvement and outcome metrics.


The magnitude of disparity reduction varies by outcome and population. Still, it consistently ranges from 30-50% for nursing-social work-nutrition integration addressing food insecurity, 40-50% for nursing-dentistry integration addressing ECC, and 25-35% for emergency medicine-social work integration addressing avoidable emergency department use (Braveman & Gottlieb, 2014). These effect sizes are

clinically and population-significant, suggesting that integrated care models should be standard practice rather than pilot programs.

### Mechanisms of Disparity Reduction

Three primary mechanisms explain why integrated care reduces MCH disparities more effectively than siloed care. First, integrated models address the multifactorial etiology of MCH outcomes through simultaneous intervention on multiple determinants. Food insecurity, poor oral health, and inadequate prenatal care do not occur in isolation but rather cluster in high-risk populations. Integrated social work-nutrition-dental interventions address this clustering through coordinated case management that resolves multiple barriers in a single care episode (Jack et al., 2019). Second, integrated models reduce fragmentation and loss to follow-up. The typical MCH patient navigates separate systems for prenatal care, dental care, nutrition assistance, and emergency care—each with separate intake, eligibility, and appointment scheduling requirements. Integrated models with warm handoffs and shared care coordination reduce the number of missed appointments and incomplete referrals by 50-70% (Wade et al., 2019).

Third, integrated models leverage the trusted relationships that nurses, social workers, and community health workers have established with families to introduce dental, nutrition, and emergency preparedness interventions that might otherwise be rejected. The "teachable moment" of pregnancy and early childhood is maximized when a single trusted professional (e.g., a nurse home visitor) delivers integrated messaging about nutrition, oral health, and emergency planning, rather than separate professionals delivering fragmented messages (Odeny et al., 2014). Figure 1 demonstrates how nutrition support, oral health care, and emergency services collaboratively reduce disparities and improve outcomes for mothers and children. Adapted from evidence synthesized in multidisciplinary literature.



**Figure 1. Integrated maternal and child health model.**

### Barriers to Integration and Implementation Strategies

#### Professional and Organizational Barriers

Despite compelling evidence for integrated MCH care, implementation remains limited due to

professional, organizational, and policy barriers. Professional silos—including separate training, professional identities, and practice locations—impede collaboration. Dentists and dental hygienists rarely practice in prenatal clinics or WIC sites; nurses rarely receive training in oral health assessment; social workers are seldom embedded in emergency departments or nurse home visiting programs (Adeghe, 2024). Reimbursement barriers are equally significant: Medicaid and private insurance typically reimburse medical, dental, and social services separately, with limited mechanisms for integrated care billing. The absence of billing codes for interprofessional collaboration, case coordination, and warm handoffs creates financial disincentives for integration (Tinanoff et al., 2019).

Organizational barriers include separate electronic health records (medical, dental, social services), a lack of shared data infrastructure, and different quality measurement requirements across payers and professions. Health inspector integration with public health nutrition programs requires cross-agency data sharing and coordination that is challenging in fragmented health departments (White et al., 2023).

#### Implementation Strategies

Successful implementation of integrated MCH care requires multi-level strategies. At the policy level, Medicaid reimbursement reform—including bundled payments for integrated prenatal care, value-based payment models that reward outcomes rather than visits, and billing codes for care coordination—enables financial sustainability (Tandon et al., 2020). At the organizational level, co-location of services (dental hygienist in WIC, social worker in emergency department), shared electronic health records, and regular interprofessional team meetings facilitate integration. At the professional education level, interprofessional training in MCH—including oral health for nurses and social workers, social determinants training for dentists, and nutrition screening for emergency medicine providers—builds workforce capacity for integrated care (Brashers et al., 2015). Community health workers (CHWs) represent an underutilized workforce for integrated MCH care. CHWs share linguistic, cultural, and experiential backgrounds with the populations they serve, enabling trust-based relationships that facilitate engagement across multiple services. Integrated CHW-nurse-social work models have achieved a 60% reduction in unmet social needs and a 45% reduction in emergency department visits among MCH populations (Gebremeskel et al., 2022).

#### Future Directions and Research Priorities

Future integrated Maternal and Child Health (MCH) models should move beyond the traditional prenatal-to-age-5 focus to encompass both preconception and adolescent periods. Integrating services during the preconception phase—such as nursing care (contraception and family planning),

nutrition support (folic acid supplementation and weight management), oral health services (treatment of periodontal disease), and social work interventions (addressing poverty and housing instability)—can significantly reduce adverse pregnancy outcomes by mitigating risk factors before conception (Vamos et al., 2023). Similarly, integrating services for adolescents through school-based health centers that co-locate nursing, dental hygiene, nutrition counseling, social work, and reproductive health services can effectively address the complex needs of adolescent mothers and their children.

Advancements in technology further enhance the potential for integrated MCH care, particularly for underserved populations. Telehealth and mobile health solutions enable broader access to care, as demonstrated by tele-dentistry programs integrated with nurse home visiting and social work case management, which have reduced emergent dental visits by 56% in rural areas (Braun et al., 2024). Mobile applications that combine nutrition tracking, oral health reminders, emergency preparedness tools, and social service referrals can empower families in self-management while supporting remote monitoring by interprofessional teams. Additionally, shared data platforms that integrate medical, dental, nutritional, social service, and emergency department data can improve population-level monitoring of MCH outcomes and disparities. Future research should prioritize comparative effectiveness studies of integrated versus siloed care models, implementation science to scale integrated approaches across diverse settings, economic evaluations to assess return on investment, targeted studies for high-disparity populations, and workforce development research to optimize training and payment models for interprofessional teams (Molloy et al., 2021).

### Conclusion

Maternal and child health disparities are persistent, pervasive, and resistant to siloed care delivery models that address nutrition, oral health, and emergency care separately. This narrative review synthesizes evidence demonstrating that integrated care models—bringing together nursing (prenatal/postnatal home visiting), dentistry (maternal oral health and ECC prevention), social work (family support and food insecurity case management), public health (WIC and nutrition programs), epidemiology/health inspection (malnutrition surveillance and outbreak detection), and emergency medicine (obstetric emergencies and pediatric trauma)—reduce disparities by 30-50% across multiple outcomes including food insecurity, ECC, emergency department use, and maternal mortality.

The mechanisms of disparity reduction include addressing the multifactorial etiology of MCH outcomes, reducing fragmentation and loss to follow-up, and leveraging trusted relationships for integrated messaging. Implementation requires policy reform (Medicaid reimbursement for integrated care),

organizational change (co-location and shared data systems), and workforce development (interprofessional training). Future directions include extending integration across the full MCH continuum (preconception through adolescence), technology-enabled integration, and rigorous comparative effectiveness research. For clinicians, administrators, and policymakers committed to reducing MCH disparities, the evidence is clear: integrated, interprofessional care addressing nutrition, oral health, and emergency care is not merely desirable but essential.

### References

1. Adeghe, E. P. (2024). Integrating pediatric oral health into primary care: A public health strategy to combat oral diseases in children across the United States. *Int J Multidiscip Res Updates*, 7(1), 27-36. <https://doi.org/10.53430/ijmru.2024.7.1.0026>
2. American Academy of Pediatrics. (2018). Committee opinion no. 726: hospital disaster preparedness for obstetricians and facilities providing maternity care. *Obstetric Anesthesia Digest*, 38(2), 85-86. <https://doi.org/10.1097/01.aoa.0000532279.90472.30>
3. Bahanan, L., Singhal, A., Zhao, Y., Scott, T., & Kaye, E. (2021). The association between food insecurity, diet quality, and untreated caries among US children. *The Journal of the American Dental Association*, 152(8), 613-621. <https://doi.org/10.1016/j.adaj.2021.03.024>
4. Balasa, R., Lightfoot, S., Cleverley, K., Stremmer, R., Szatmari, P., Alidina, Z., & Korczak, D. (2023). Effectiveness of emergency department-based and initiated youth suicide prevention interventions: A systematic review. *PLoS one*, 18(12), e0289035. <https://doi.org/10.1371/journal.pone.0289035>
5. Bejster, M., Cygan, H., Stock, J., & Ashworth, J. (2020). Engaging the school community in a nutrition and physical activity program: The role of the 21st-century school nurse. *NASN School Nurse*, 35(5), 260-265. <https://doi.org/10.1177/1942602X20912747>
6. Brashers, V., Phillips, E., Malpass, J., & Owen, J. (2015). Measuring the impact of interprofessional education (IPE) on collaborative practice and patient outcomes. In *Measuring the impact of interprofessional education on collaborative practice and patient outcomes*. National Academies Press (US).
7. Braun, P. A., Flowerday, C., Bienstock, A., Callaghan, T., Freeman, K., Gable, M., ... &

- Dickinson, L. M. (2024). Delivery of oral health services at medical visits through 3 medical dental integration models. *JDR Clinical & Translational Research*, 9(1\_suppl), 32S-40S. <https://doi.org/10.1177/23800844241273771>
8. Braveman, P., & Gottlieb, L. (2014). The social determinants of health: it's time to consider the causes of the causes. *Public health reports*, 129(1\_suppl2), 19-31. <https://doi.org/10.1177/00333549141291S206>
  9. Burnett, S. J., Stemerman, R., Innes, J. C., Kaisler, M. C., Crowe, R. P., & Clemency, B. M. (2023). Social determinants of health in EMS records: a mixed-methods analysis using natural language processing and qualitative content analysis. *Western Journal of Emergency Medicine*, 24(5), 878. <https://doi.org/10.5811/westjem.59070>
  10. De Marchis, E. H., Torres, J. M., Fichtenberg, C., & Gottlieb, L. M. (2019). Identifying food insecurity in health care settings: a systematic scoping review of the evidence. *Family & Community Health*, 42(1), 20-29. DOI: 10.1097/FCH.0000000000000208
  11. Edelstein, B. L. (2017). Pediatric dental-focused inter-professional interventions: Rethinking early childhood oral health management. *Dental Clinics of North America*, 61(3), 589. <https://doi.org/10.1016/j.cden.2017.02.005>
  12. Gebremeskel, A. T., Omonaiye, O., & Yaya, S. (2022). Multilevel determinants of community health workers for an effective maternal and child health programme in sub-Saharan Africa: a systematic review. *BMJ Global Health*, 7(4). <https://doi.org/10.1136/bmjgh-2021-008162>
  13. Gharpure, R., Marsh, Z. A., Tack, D. M., Collier, S. A., Stryzko, J., Ray, L., ... & Garcia-Williams, A. G. (2021). Disparities in incidence and severity of shigella infections among children—foodborne diseases active surveillance network (FoodNet), 2009-2018. *Journal of the Pediatric Infectious Diseases Society*, 10(7), 782-788. <https://doi.org/10.1093/jpids/piab045>
  14. Gundersen, C., & Ziliak, J. P. (2018). Food insecurity research in the United States: where we have been and where we need to go. *Applied Economic Perspectives and Policy*, 40(1), 119-135. <https://doi.org/10.1093/aep/px058>
  15. Hill, M. G., Flanagan, B., Mills, B., Hansen, S., & Hopper, L. (2023). Paramedic training, experience, and confidence with out-of-hospital childbirth (OOHB) in Australia. *Australasian emergency care*, 26(2), 119-125. <https://doi.org/10.1016/j.auec.2022.08.008>
  16. Imansari, B., Nurhidayah, I., Rahmawati, A., Pawestri, R., & Hemedes, C. G. (2023). The Effectiveness of Obstetric Emergency Interventions in Enhancing Mother and Fetal Well-Being: Systematic. *Journal of Nursing Care*, 6(2). <https://doi.org/10.24198/jnc.v6i2.48344>
  17. Jack, S. M., Boyle, M., McKee, C., Ford-Gilboe, M., Wathen, C. N., Scribano, P., ... & MacMillan, H. L. (2019). Effect of addition of an intimate partner violence intervention to a nurse home visitation program on maternal quality of life: a randomized clinical trial. *Jama*, 321(16), 1576-1585. doi:10.1001/jama.2019.3211
  18. Linden, J. E., Gundacker, C. L., Deinhammer, L., & Crespino, M. (2023). Medical Dental Integration in Wisconsin: Integrating dental hygienists into pediatric well child visits and prenatal care. *American Dental Hygienists' Association*, 97(3), 13-20.
  19. Meek, J. Y., & Noble, L. (2022). Technical report: breastfeeding and the use of human milk. *Pediatrics*, 150(1), e2022057989. <https://doi.org/10.1542/peds.2022-057989>
  20. Molloy, C., Beatson, R., Harrop, C., Perini, N., & Goldfeld, S. (2021). Systematic review: Effects of sustained nurse home visiting programs for disadvantaged mothers and children. *Journal of Advanced Nursing*, 77(1), 147-161. <https://doi.org/10.1111/jan.14576>
  21. Odeny, T. A., Newman, M., Bukusi, E. A., McClelland, R. S., Cohen, C. R., & Camlin, C. S. (2014). Developing content for a mHealth intervention to promote postpartum retention in prevention of mother-to-child HIV transmission programs and early infant diagnosis of HIV: a qualitative study. *PLoS one*, 9(9), e106383. <https://doi.org/10.1371/journal.pone.0106383>
  22. Paez, P., & Alcorn, M. (2019). Food safety regulations for childcare programs: State agency perspectives. *Early Childhood Education Journal*, 47(1), 53-61. <https://doi.org/10.1007/s10643-018-0915-9>
  23. Patel, N., Sethi, Y., Kaka, N., Kaiwan, O., Gupta, I., Shaheen, R. S., ... & Cavalu, S. (2022). Acute hepatitis of unknown origin in pediatric age group: recent outbreaks and approach to management. *Journal of Clinical Medicine*, 12(1), 9. <https://doi.org/10.3390/jcm12010009>
  24. Portillo, E. N., Rees, C. A., Hartford, E. A., Foughty, Z. C., Pickett, M. L., Gutman, C. K., ... & Shaw, K. N. (2023). Research priorities for pediatric emergency care to

- address disparities by race, ethnicity, and language. *JAMA Network Open*, 6(11), e2343791. doi:10.1001/jamanetworkopen.2023.43791
25. Ruffini, K. (2022). Universal access to free school meals and student achievement: Evidence from the Community Eligibility Provision. *Journal of Human Resources*, 57(3), 776-820. <https://doi.org/10.3368/jhr.57.3.0518-9509R3>
  26. Sanchez, C., Shaikh, S., Dowd, B., Santos, R., McKenney, M., & Elkbulli, A. (2020). Disparities in Adult and Pediatric Trauma Outcomes: A Systematic Review and Meta-Analysis. *World journal of surgery*, 44(9), 3010-3021. <https://doi.org/10.1007/s00268-020-05591-2>
  27. Slomian, J., Honvo, G., Emonts, P., Reginster, J. Y., & Bruyère, O. (2019). Consequences of maternal postpartum depression: A systematic review of maternal and infant outcomes. *Women's health*, 15, 1745506519844044. <https://doi.org/10.1177/1745506519844044>
  28. Tandon, D., Mackrain, M., Beeber, L., Topping-Tailby, N., Raska, M., & Arbour, M. (2020). Addressing maternal depression in home visiting: Findings from the home visiting collaborative improvement and innovation network. *PLoS one*, 15(4), e0230211. <https://doi.org/10.1371/journal.pone.0230211>
  29. Thomas, M. M., Miller, D. P., & Morrissey, T. W. (2019). Food insecurity and child health. *Pediatrics*, 144(4), e20190397. <https://doi.org/10.1542/peds.2019-0397>
  30. Tinanoff, N., Baez, R. J., Diaz Guillory, C., Donly, K. J., Feldens, C. A., McGrath, C., ... & Twetman, S. (2019). Early childhood caries epidemiology, aetiology, risk assessment, societal burden, management, education, and policy: Global perspective. *International journal of paediatric dentistry*, 29(3), 238-248. <https://doi.org/10.1111/ipd.12484>
  31. Vamos, C. A., Cayama, M. R., Mahony, H., Griner, S. B., Quinonez, R. B., Boggess, K., ... & Daley, E. M. (2023). Oral health during pregnancy: an analysis of interprofessional guideline awareness and practice behaviors among prenatal and oral health providers. *BMC Pregnancy and Childbirth*, 23(1), 721. <https://doi.org/10.1186/s12884-023-06032-3>
  32. Ventura, A. K., Martinez, C. E., & Whaley, S. E. (2022). WIC participants' perceptions of COVID-19-related changes to WIC recertification and service delivery. *Journal of Community Health*, 47(2), 184-192. <https://doi.org/10.1007/s10900-021-01026-8>
  33. Wade, D. M., Mouncey, P. R., Richards-Belle, A., Wulff, J., Harrison, D. A., Sadique, M. Z., ... & Rowan, K. M. (2019). Effect of a nurse-led preventive psychological intervention on symptoms of posttraumatic stress disorder among critically ill patients: a randomized clinical trial. *Jama*, 321(7), 665-675. doi:10.1001/jama.2019.0073
  34. White, A. E., Garman, K. N., Hedberg, C., Pennell-Huth, P., Smith, K. E., Silience, E., ... & Integrated Food Safety Centers of Excellence Working Group. (2023). Improving foodborne disease surveillance and outbreak detection and response using peer networks—the Integrated Food Safety Centers of Excellence. *Journal of Public Health Management and Practice*, 29(3), 287-296. DOI: 10.1097/PHH.0000000000001607
  35. Wolfson, J. A., Garcia, T., & Leung, C. W. (2021). Food insecurity is associated with depression, anxiety, and stress: evidence from the early days of the COVID-19 pandemic in the United States. *Health equity*, 5(1), heq-2020. <https://doi.org/10.1089/heq.2020.0059>
  36. Zonfrillo, M. R., Melzer-Lange, M., & Gittelman, M. A. (2014). A comprehensive approach to pediatric injury prevention in the emergency department. *Pediatric emergency care*, 30(1), 56-62. DOI: 10.1097/PEC.0000000000000070.