Saudi Journal of Medicine and Public Health

https://saudijmph.com/index.php/pub

https://doi.org/10.64483/202631855

Infant Apnea: Etiology, Pathophysiology, Clinical Manifestations, Diagnostic Evaluation, Nursing Assessment, and Evidence-Based Management Strategies.

Ibtisam Mohammed Alanazi[(1)] , Nawal Mohammed Hamad Arishi[(2)] , Fatimah Hussin Najmi[(3)] , Mohammed Abdullah Yahya Asiri[(4)] , Amraa Hamzah A. Alhawsah[(5)] , Khaled Abdullah Shafi Al-Toum[(6)] , Reem Falah Almutairi[(7)] , Nuha Fahad Almutlaq[(8)] , Najah Shalil Alanazi[(9)] , Nada Fahad Mutlaq Almutlaq[(8)]

(1)Maternity and Children Hospital, Hafar Al-Batin, Saudi Arabia, Ministry of Health, Saudi Arabia
(2)Jazan Health Cluster, Jazan, Saudi Arabia, Ministry of Health, Saudi Arabia
(3)Samtah General Hospital, Jazan, Saudi Arabia, Ministry of Health, Saudi Arabia
(4)Abha Prison Health Center, Abha, Saudi Arabia, Ministry of Interior, Saudi Arabia
(5)King Fahd Hospital, Madinah, Saudi Arabia, Ministry of Health, Saudi Arabia
(6)Sajer General Hospital, Saudi Arabia, Ministry of Health, Saudi Arabia
(7)Dharma Health Center, Saudi Arabia, Ministry of Health, Saudi Arabia
(8)Imam Abdulrahman Al Faisal Hospital, Riyadh, Saudi Arabia, Ministry of Health, Saudi Arabia
(9)Uhud Hospital, Madinah, Saudi Arabia, Ministry of Health, Saudi Arabia

Abstract

Background: Infant apnea is a clinical condition characterized by the cessation or significant reduction of breathing. It may occur in premature and term infants and can result from respiratory, neurological, cardiovascular, infectious, and metabolic disorders. Early recognition and appropriate management are essential to prevent complications, particularly in vulnerable neonates.

Aim: This review aims to examine the etiology, pathophysiology, clinical manifestations, diagnostic evaluation, differential diagnosis, treatment, and nursing management of infant apnea, with emphasis on multidisciplinary care and parental education.

Methods: A narrative review approach was used to discuss the available clinical literature and established recommendations concerning infant apnea. Relevant topics included risk factors, underlying mechanisms, clinical assessment, diagnostic investigations, therapeutic interventions, specialist consultations, and strategies for improving healthcare team outcomes.

Results: Infant apnea has multiple etiologies, including apnea of prematurity, respiratory infections, airway abnormalities, cardiac disorders, neurological conditions, metabolic disturbances, and sepsis. Clinical manifestations include breathing pauses, bradycardia, oxygen desaturation, cyanosis, pallor, and altered responsiveness. Diagnostic evaluation depends on the clinical history and physical examination, with laboratory tests, electrocardiography, neuroimaging, and other investigations selected according to clinical indications. Management focuses on stabilization, identification of the underlying cause, respiratory support, and targeted treatment. Caffeine therapy and continuous positive airway pressure are important interventions for selected premature infants. Nursing surveillance, safe sleep education, and caregiver training remain essential components of care.

Conclusion: Early diagnosis, appropriate treatment, continuous monitoring, and coordinated multidisciplinary management are necessary to reduce complications and improve outcomes in infants with apnea.

Keywords: Infant apnea, apnea of prematurity, neonatal respiratory disorders, cardiorespiratory monitoring, caffeine therapy, nursing management, brief resolved unexplained event.

Introduction

Apnea in infants is characterized by temporary cessation or significant interruption of breathing and represents an important clinical finding during the neonatal and early infancy periods. It may result from physiological immaturity, developmental abnormalities in respiratory control, or underlying pathological conditions affecting the respiratory, cardiovascular, neurological, or metabolic systems. Although some breathing irregularities are considered normal during early infancy, prolonged or clinically significant respiratory pauses require careful assessment to identify their underlying causes and prevent potential complications. One common physiological breathing pattern observed in young infants is periodic breathing. This phenomenon consists of short respiratory pauses, usually lasting approximately 5 to 10 seconds, followed by periods of regular breathing. Periodic breathing is generally considered a normal developmental pattern and does

*Corresponding author bsoom09@icloud.com (Ibtisam Mohammed Alanazi).
Receive Date: 5 September 2026, Revise Date: 30 October 2026, Accept Date: 16 November 2026

not usually indicate an underlying respiratory disorder. It occurs most frequently between 2 and 4 weeks of age and typically resolves spontaneously by approximately 6 months. Unlike pathological apnea, periodic breathing is not usually associated with significant oxygen desaturation, bradycardia, or changes in skin color or muscle tone. Nevertheless, distinguishing physiological respiratory irregularities from clinically significant apnea is essential, particularly in premature infants and neonates with associated medical conditions.[1,2]

Apnea occurs more frequently in premature infants because their respiratory control mechanisms and neurological pathways responsible for maintaining regular ventilation are not fully developed. Apnea of prematurity is defined as an episode of cessation of breathing lasting at least 20 seconds or a shorter respiratory pause accompanied by bradycardia, oxygen desaturation, or cyanosis in an infant born before 37 weeks of gestation. The condition is associated with immaturity of the central nervous system, reduced responsiveness to changes in carbon dioxide and oxygen concentrations, and inadequate coordination of respiratory activity. Its frequency and severity are generally inversely related to gestational age and birth weight. In full-term infants, unexplained episodes of respiratory cessation are classified as apnea of infancy. This condition involves an interruption of breathing lasting 20 seconds or longer or a shorter pause accompanied by clinically significant manifestations, including bradycardia, cyanosis, pallor, or marked hypotonia. Such episodes require clinical evaluation to exclude respiratory infections, airway abnormalities, neurological disorders, cardiac conditions, metabolic disturbances, and other underlying diseases. Based on the physiological mechanism, infant apnea is classified into three main types: central, obstructive, and mixed apnea. Central apnea occurs when respiratory activity from the central nervous system temporarily stops, resulting in the absence of both airflow and respiratory effort. Obstructive apnea develops when airflow is interrupted by partial or complete upper airway obstruction despite continued respiratory effort. Mixed apnea combines both mechanisms, beginning with a central respiratory pause and followed by continued respiratory effort against an obstructed airway. Mixed apnea represents the most frequently observed pattern among premature infants. Understanding the definition, classification, and physiological mechanisms of infant apnea is essential for appropriate clinical assessment and management. Early recognition, accurate differentiation between physiological and pathological respiratory pauses, and identification of associated risk factors help guide monitoring and treatment strategies and reduce the risk of complications related to recurrent hypoxemia and bradycardia.

Etiology

The etiology of infant apnea is multifactorial and includes a wide range of physiological, developmental, neurological, respiratory, infectious, metabolic, and structural conditions. The underlying causes vary according to gestational age, postnatal age, clinical presentation, and the type of apnea. Identifying the specific etiology is essential for determining appropriate treatment, preventing recurrent episodes, and reducing the risk of complications associated with prolonged respiratory interruption. Apnea is particularly common among premature infants because of the incomplete maturation of the neurological and respiratory systems responsible for maintaining regular breathing.[3–5] Apnea of prematurity represents one of the most frequent causes of respiratory pauses in neonatal intensive care units. It occurs predominantly in infants born before 37 weeks of gestation, particularly those born before 28 weeks. The primary mechanism involves immaturity of the central respiratory control centers, reduced responsiveness to changes in arterial oxygen and carbon dioxide concentrations, and inadequate coordination of respiratory muscle activity. Premature infants may also demonstrate an immature response to airway stimulation and altered regulation of upper airway muscle tone. These developmental limitations increase susceptibility to recurrent central, obstructive, and mixed apnea.




Figure-1: Etiology of Infant Apnea.

Feeding-related respiratory disturbances may also contribute to apnea episodes in premature infants. During feeding, immature coordination between sucking, swallowing, and breathing can interrupt normal ventilation and produce oxygen desaturation. An immature laryngeal chemoreflex may further contribute to respiratory inhibition following exposure of the laryngeal structures to liquids or other stimuli. Respiratory deterioration after feeding may also be associated with diaphragmatic fatigue and reduced respiratory reserve. Although gastroesophageal reflux is frequently suspected as a cause of apnea, available evidence indicates that reflux is not generally responsible for most apnea episodes in infants.[6] In newborns, apnea occurring shortly after delivery may result from several perinatal and maternal factors. These include birth asphyxia, maternal exposure to

medications or substances that depress respiratory activity, neonatal infections, metabolic abnormalities, and congenital structural or neurological disorders. Birth asphyxia may impair respiratory control through hypoxic and ischemic injury to the central nervous system. Maternal administration of opioids, sedatives, anesthetic agents, or other respiratory depressants may also temporarily affect neonatal respiratory function.

Central apnea is associated with conditions that interfere with the activity of the brainstem respiratory centers or neural pathways responsible for respiratory regulation. Important causes include central nervous system infections, particularly meningitis and encephalitis, as well as head trauma resulting from birth injury or abusive trauma. Exposure to toxins and certain medications may suppress respiratory drive. Infectious conditions such as pertussis and infant botulism can also produce apnea through respiratory center dysfunction or neuromuscular impairment. In addition, inherited metabolic disorders, including mitochondrial diseases, Pompe disease, Leigh syndrome, and mucopolysaccharidoses, may affect neurological and respiratory function. Metabolic disturbances such as hypoglycemia, hypocalcemia, and metabolic acidosis can further interfere with normal respiratory regulation. Congenital conditions, including congenital central hypoventilation syndrome, Down syndrome, and Arnold-Chiari malformation, are additional recognized causes of central apnea. Obstructive apnea develops when airflow becomes partially or completely restricted despite continued respiratory effort. Potential causes include obstructive sleep apnea, respiratory infections such as pneumonia and croup, vocal cord paralysis, and congenital abnormalities affecting the upper airway. Structural conditions such as Pierre Robin sequence may produce airway narrowing and interfere with normal ventilation, particularly during sleep. Mixed apnea involves the combination of central respiratory inhibition and upper airway obstruction. It occurs predominantly in premature infants because of the simultaneous immaturity of central respiratory control and upper airway stabilization mechanisms. However, mixed apnea may also occur in association with gastroesophageal reflux, pertussis, and bronchiolitis. These conditions may affect respiratory drive, airway patency, or both mechanisms. From a nursing perspective, understanding the diverse causes of infant apnea is essential for identifying high-risk neonates, recognizing associated clinical manifestations, assessing maternal and perinatal history, and monitoring respiratory patterns. Careful assessment of gestational age, feeding history, medication exposure, infection indicators, and neurological status helps nurses identify potential contributing factors and communicate relevant findings promptly to the healthcare team [7].

Epidemiology

The epidemiology of infant apnea varies considerably according to gestational age, birth weight, postnatal development, and the presence of underlying medical conditions. The exact prevalence and incidence of infant apnea remain uncertain because of differences in diagnostic definitions, monitoring techniques, study populations, and the clinical criteria used to identify respiratory pauses. Furthermore, some episodes of apnea occur without noticeable clinical manifestations and may only be detected through continuous cardiorespiratory monitoring in neonatal intensive care units. Despite these limitations, available evidence indicates that apnea is substantially more common among premature infants than among full-term newborns. Apnea of prematurity represents the most frequently encountered form of infant apnea, particularly among neonates born at very early gestational ages. Its incidence is inversely proportional to gestational age, meaning that the likelihood of developing apnea increases as gestational age decreases. Nearly all infants born before 28 weeks of gestation experience at least one episode of apnea during the neonatal period. Among infants born between 33 and 34 weeks of gestation, approximately 50% experience an episode, while the incidence decreases to approximately 20% by 34 weeks of gestation.[8] This relationship reflects the progressive maturation of the central nervous system, respiratory control mechanisms, and respiratory muscle coordination as pregnancy advances.

The high incidence among extremely premature infants has important implications for neonatal healthcare services. These infants frequently require admission to neonatal intensive care units, where continuous monitoring of respiratory rate, oxygen saturation, and heart rate is necessary to identify episodes of apnea and associated bradycardia or oxygen desaturation. The frequency of episodes generally decreases with advancing postnatal and postmenstrual age as respiratory control mechanisms mature. However, the time required for complete resolution varies among infants and may be influenced by associated complications, including respiratory distress syndrome, bronchopulmonary dysplasia, infection, and neurological immaturity. In contrast, apnea is relatively uncommon among full-term infants. The estimated incidence in this population is approximately 1 case per 1,000 infants.[9] When apnea occurs in full-term newborns, it is particularly important to investigate potential underlying pathological conditions rather than automatically attributing the episodes to developmental immaturity. Respiratory infections, neurological abnormalities, metabolic disturbances, congenital airway disorders, and cardiovascular conditions may contribute to its occurrence. From a nursing and clinical perspective, understanding the

epidemiological distribution of infant apnea helps identify neonates who require closer observation and specialized care. Premature infants, particularly those born before 28 weeks of gestation, represent the highest-risk population. Appropriate risk assessment, continuous cardiorespiratory monitoring, and timely recognition of respiratory abnormalities are essential components of neonatal nursing practice. These measures support early intervention, reduce the likelihood of complications, and improve the quality and safety of neonatal care [8][9].

Pathophysiology

The pathophysiology of infant apnea is complex and involves interactions between the developing central nervous system, respiratory control mechanisms, upper airway structures, respiratory muscles, and environmental or physiological stressors. The condition occurs more frequently in premature infants because the neurological pathways responsible for regulating breathing have not reached functional maturity. The respiratory centers located in the brainstem play a central role in maintaining regular ventilation through the integration of chemical, mechanical, and neurological signals. In premature neonates, these regulatory mechanisms demonstrate reduced responsiveness and inadequate coordination, making respiratory activity vulnerable to temporary interruption in response to internal and external stimuli.[10,11] One of the major mechanisms contributing to apnea involves the immature ventilatory response to changes in blood oxygen and carbon dioxide concentrations. In adults and older children, hypoxia generally stimulates increased respiratory activity, while hypercarbia enhances ventilation to eliminate excess carbon dioxide. However, neonates, particularly premature infants, may demonstrate a different response. Initial exposure to hypoxia or hypercarbia may produce a brief increase in respiratory rate, followed by respiratory depression and, in some cases, complete cessation of breathing. This abnormal response reflects the immaturity of central and peripheral chemoreceptors and their associated neural pathways. Consequently, premature infants may experience recurrent respiratory pauses when exposed to physiological disturbances that would normally stimulate ventilation in older individuals.

Feeding-related apnea represents another important pathophysiological mechanism. Effective feeding requires precise coordination between sucking, swallowing, and breathing. In premature infants, incomplete neurological development may interfere with this coordination, resulting in reduced ventilation, oxygen desaturation, and respiratory pauses. Furthermore, stimulation of laryngeal sensory receptors by liquids or other substances may activate an exaggerated laryngeal chemoreflex. This reflex can produce protective airway responses, including swallowing, coughing, laryngeal closure, and inhibition of respiratory activity. Although these responses help prevent aspiration, their exaggerated activation in immature infants may contribute to apnea and bradycardia. Several systemic and environmental factors can further impair respiratory regulation. Anemia reduces the oxygen-carrying capacity of the blood, while hypoglycemia interferes with cellular energy metabolism and normal neurological activity. Hypothermia increases metabolic stress and may adversely affect respiratory function. Exposure to medications, toxins, or substances that depress the central nervous system can also reduce respiratory drive. These conditions may independently contribute to apnea or aggravate pre-existing respiratory instability in premature neonates.

Thoracic and respiratory muscle characteristics also influence the development of infant apnea. The neonatal thoracic cage is highly compliant because its skeletal structures are incompletely developed. This increased flexibility makes the chest wall more susceptible to inward movement during inspiration, particularly when negative intrathoracic pressure increases. Consequently, premature infants must generate greater respiratory effort to maintain adequate ventilation. Increased work of breathing may lead to respiratory muscle fatigue, reduced tidal volume, inadequate alveolar ventilation, and progressive respiratory compromise. In severe cases, these mechanisms may contribute to respiratory failure and recurrent apnea. The pathophysiology of apnea in full-term infants differs from that observed in premature neonates. Apnea of infancy generally results from identifiable underlying pathological conditions rather than isolated developmental immaturity. The mechanisms vary according to the responsible disorder and may involve abnormal airway protection, upper airway obstruction, neurological dysfunction, or disturbances in respiratory regulation. Gastroesophageal reflux disease has been reported as an associated condition in approximately 48% of cases, upper airway abnormalities and obstruction in 37%, and neurological diseases in 19%. However, these conditions may overlap, and individual infants may have multiple contributing factors.[12]

In central apnea, suppression or interruption of brainstem respiratory output results in the absence of respiratory effort and airflow. In obstructive apnea, respiratory drive remains active, but airflow becomes restricted because of partial or complete upper airway obstruction. Mixed apnea combines both mechanisms, with an initial period of absent respiratory effort followed by continued respiratory attempts against an obstructed airway. These mechanisms may produce intermittent hypoxemia, carbon dioxide retention, and secondary cardiovascular responses. From a nursing perspective, understanding the pathophysiological

mechanisms of infant apnea is essential for interpreting changes in respiratory patterns, oxygen saturation, and heart rate. Continuous assessment of respiratory effort, chest movement, airway patency, and associated physiological abnormalities helps nurses recognize different apnea patterns and identify deterioration. Early recognition and appropriate intervention are important for maintaining adequate ventilation, preventing recurrent hypoxemia, and reducing complications associated with prolonged or repeated respiratory interruption.




Figure-2: Pathophysiology of infant Apnea.

History and Physical Examination

The history and physical examination represent essential components of the initial assessment of infants presenting with suspected apnea. A systematic clinical evaluation helps distinguish true apneic episodes from normal physiological breathing variations, identify potential underlying causes, determine the severity of the condition, and guide appropriate diagnostic and therapeutic interventions. The initial assessment should establish whether the reported event represents true cessation of breathing, periodic breathing, or a breath-holding spell. Periodic breathing is characterized by brief respiratory pauses that are generally not associated with significant oxygen desaturation, bradycardia, or changes in skin color. In contrast, pathological apnea may involve prolonged respiratory interruption accompanied by cyanosis, pallor, reduced muscle tone, bradycardia, or oxygen desaturation. Following confirmation of a suspected apneic event, the clinical approach should consider the infant's gestational age at birth. This distinction is important because apnea of prematurity generally reflects developmental immaturity of respiratory control, whereas apnea in full-term infants requires careful investigation for underlying pathological conditions.

History Taking in Premature Infants

In infants born before 37 weeks of gestation, the assessment should focus on identifying characteristics consistent with apnea of prematurity and determining whether additional medical conditions contribute to respiratory instability. Apnea of prematurity is associated with immature central respiratory regulation and generally resolves as the infant approaches approximately 40 weeks of postmenstrual age, although resolution may occur later in some infants.[13–15] Important historical information includes gestational age at birth, birth weight, postnatal age, previous episodes of apnea, frequency and duration of respiratory pauses, and any interventions required during previous events. Nurses and healthcare professionals should also establish whether episodes occur during sleep, wakefulness, feeding, or changes in body position. Particular attention should be given to events associated with feeding because immature coordination between sucking, swallowing, and breathing may contribute to respiratory disturbances. The assessment should document associated clinical manifestations, including bradycardia, oxygen desaturation, cyanosis, pallor, and changes in muscle tone. Information regarding respiratory support, medication administration, feeding tolerance, and previous neonatal complications is also important. The physical examination should evaluate general appearance, respiratory effort, chest movement, airway patency, cardiovascular status, neurological responses, and signs of developmental maturity. Healthcare professionals should also assess for conditions that may aggravate apnea, including infection, anemia, hypoglycemia, hypothermia, and electrolyte disturbances.[13–15]

History Taking in Full-Term Infants

In full-term infants born at or beyond 37 weeks of gestation, episodes of apnea require a broader diagnostic assessment because they are more likely to be associated with identifiable pathological conditions. Important reported causes include gastroesophageal reflux disease, upper airway abnormalities or obstruction, and neurological disorders, accounting for approximately 48%, 37%, and 19% of cases, respectively. These conditions may coexist in individual infants.[12] The history should explore the circumstances surrounding each episode, including its onset, duration, frequency, severity, and relationship to feeding or sleeping. Feeding-related symptoms require particular attention, including choking, coughing, regurgitation, vomiting, difficulty swallowing, and poor feeding tolerance. These findings may suggest swallowing dysfunction, aspiration, or associated gastrointestinal problems. Assessment of upper airway symptoms should include questions about stridor, noisy breathing, snoring, nasal congestion, recurrent respiratory infections, and breathing difficulties associated with particular sleeping positions. Parents should also be asked whether respiratory symptoms improve or worsen when the infant's head or body position changes. Neurological history should explore abnormal movements, unusual muscle tone, reduced responsiveness, seizures, developmental abnormalities, and episodes of altered consciousness.[12,16,17]

Physical Examination

The physical examination of full-term infants should follow a systematic approach covering respiratory, cardiovascular, neurological, and anatomical findings. Initial assessment includes respiratory rate, oxygen saturation, heart rate, temperature, skin color, level of consciousness, and general activity. Observation of respiratory patterns is important for identifying pauses, irregular breathing, increased respiratory effort, chest retractions, and abnormal respiratory sounds. The upper airway should be examined for nasal obstruction, abnormal oropharyngeal anatomy, enlarged structures, and congenital abnormalities that may interfere with airflow. Particular attention should be given to conditions associated with upper airway obstruction, including micrognathia and structural abnormalities of the palate. Respiratory assessment should include chest expansion, bilateral air entry, breath sounds, and evidence of respiratory distress. Neurological examination should evaluate muscle tone, primitive reflexes, spontaneous movements, responsiveness, and age-appropriate neurological function. Cardiovascular assessment should identify abnormal heart rates, perfusion changes, and associated signs of circulatory compromise. When clinically indicated, respiratory patterns should also be observed in different positions to identify positional airway obstruction or changes in ventilation.[12,16,17]

Nursing Considerations

Nurses play an important role in obtaining an accurate history, documenting apnea episodes, and identifying clinical deterioration. Information provided by parents and caregivers should be incorporated into the assessment, particularly regarding the timing, duration, associated manifestations, and possible triggers of respiratory pauses. Continuous observation of respiratory effort, oxygen saturation, and heart rate is essential in infants at increased risk. Accurate documentation and prompt communication of abnormal findings support early diagnosis and appropriate management. Differentiating apnea of prematurity from apnea associated with underlying pathological conditions is particularly important because the diagnostic investigations, monitoring requirements, and treatment strategies differ between these populations.

Evaluation

The diagnostic evaluation of infant apnea aims to identify the underlying cause, assess the severity of respiratory disturbances, detect associated complications, and determine the need for appropriate medical intervention. The selection of diagnostic investigations should be guided by the infant's clinical history, gestational age, physical examination findings, and characteristics of the reported apneic episodes. A comprehensive approach is particularly important because infant apnea may result from neurological, respiratory, cardiovascular, infectious, metabolic, or structural abnormalities. Diagnostic testing should therefore be individualized rather than applying the same investigations to all infants presenting with respiratory pauses. Laboratory investigations may be considered in neonates when clinical findings suggest an underlying metabolic or hematological abnormality. A complete blood count (CBC) helps identify anemia, which may reduce oxygen delivery to tissues, as well as leukocytosis or other abnormalities that may indicate infection. Measurement of serum glucose is important because hypoglycemia can interfere with normal neurological activity and respiratory regulation. Serum calcium and electrolyte measurements may also be indicated to identify metabolic disturbances that can affect neuromuscular function and respiratory control. Depending on the clinical presentation, additional biochemical investigations may be required to evaluate acid-base disturbances and other metabolic abnormalities. Infections represent important potential causes of apnea, particularly in neonates and young infants. When an infant presents with fever, hypothermia, lethargy, poor feeding, respiratory instability, or other findings suggestive of serious infection, an appropriate infectious evaluation should be performed. This may include blood cultures, urine cultures, and cerebrospinal fluid examination and culture when clinically indicated. These investigations help identify systemic bacterial infections, urinary tract infections, meningitis, and other infectious conditions that may interfere with normal respiratory function.[18]

Cardiac evaluation may also be necessary when the clinical presentation suggests an underlying cardiovascular disorder. An electrocardiogram (ECG) can help identify cardiac rhythm abnormalities associated with episodes of apparent apnea, bradycardia, or transient loss of responsiveness. Particular attention should be given to prolonged QT intervals because congenital long QT syndrome may predispose infants to potentially serious cardiac dysrhythmias and episodes of sudden clinical deterioration.[19] Neurological investigations are not routinely required in every infant with apnea. Neuroimaging, including cranial ultrasonography, computed tomography, or magnetic resonance imaging, may be considered when the history or examination suggests structural abnormalities, intracranial injury, or neurological disease. Similarly, electroencephalography (EEG) may be appropriate when episodes are associated with abnormal movements, altered consciousness, or suspected seizure activity. Consultation with pediatric neurology, cardiology, respiratory medicine, or other specialists should be guided by specific clinical findings rather than performed routinely. For infants presenting with a brief resolved unexplained event (BRUE), risk stratification is an important component of diagnostic evaluation. When the infant meets established low-risk criteria, extensive laboratory testing, imaging, and hospital investigations are generally unnecessary. Management should instead

focus on appropriate clinical assessment, caregiver education, and follow-up when indicated.[20,21] From a nursing perspective, accurate preparation for diagnostic procedures, monitoring of vital signs, observation of respiratory patterns, and documentation of abnormal findings are essential. Nurses should also communicate changes in oxygen saturation, heart rate, temperature, feeding tolerance, and neurological responsiveness to the healthcare team. These measures support timely identification of underlying conditions and contribute to safe, evidence-based management of infants presenting with apnea.

Treatment / Management

The initial step in managing infant apnea is to assess the infant's clinical condition and determine whether immediate resuscitation or stabilization is required. Evaluation should focus on airway patency, respiratory effort, heart rate, oxygen saturation, and level of responsiveness. If the infant demonstrates inadequate breathing, bradycardia, cyanosis, or significant oxygen desaturation, prompt intervention is necessary. Supportive measures may include positioning the airway, providing tactile stimulation, administering supplemental oxygen, and initiating assisted ventilation when indicated. The choice of intervention depends on the severity of the episode and the infant's response to initial measures. Following stabilization, management should focus on identifying the underlying cause and providing targeted treatment. Potential causes include prematurity, respiratory infections, metabolic disturbances, neurological disorders, cardiac abnormalities, airway obstruction, and feeding-related complications. Treatment should address the specific condition rather than apnea alone. For example, suspected infection requires appropriate antimicrobial therapy, while hypoglycemia and electrolyte disturbances require correction of the underlying metabolic abnormalities. Infants with suspected cardiac or neurological causes may require additional investigations and specialist assessment. Neonates with clinically significant apnea generally require observation in a neonatal intensive care unit or an appropriately equipped neonatal care setting. Continuous cardiorespiratory monitoring and pulse oximetry help detect recurrent episodes, bradycardia, and oxygen desaturation. Monitoring also allows healthcare professionals to assess the frequency, duration, and severity of apneic events and evaluate the infant's response to treatment. The duration of observation should depend on the clinical presentation, underlying diagnosis, and recurrence of episodes.

In infants with apnea of prematurity, treatment is considered when episodes are frequent, prolonged, require repeated stimulation, or are associated with bradycardia and hypoxemia. Nasal continuous positive airway pressure (CPAP) may be used to maintain airway patency, support lung expansion, and reduce the frequency of apnea. Methylxanthines, particularly caffeine, are also used to stimulate respiratory activity and improve respiratory drive in premature infants. These interventions are selected according to the infant's gestational age, clinical condition, and response to treatment. Continuous monitoring remains important during therapy to evaluate effectiveness and identify possible adverse effects. Methylxanthine therapy and CPAP are not routinely indicated for term infants with apnea. In these infants, management should be guided by the identified cause and the clinical severity of the episodes. Infants with feeding-related apnea may benefit from assessment by speech-language or occupational therapists. These professionals can evaluate swallowing coordination, feeding technique, and the risk of aspiration. Adjustments to feeding position, flow rate, or feeding method may be recommended when clinically appropriate. For infants classified as having a low-risk brief resolved unexplained event (BRUE), management generally involves parental reassurance and education rather than extensive diagnostic testing or treatment. Parents should receive clear information about the nature of the event, the reassuring features of the assessment, and the circumstances that require medical attention. Education may include recognizing abnormal breathing, changes in skin color, reduced responsiveness, and recurrent episodes. Caregivers should also receive guidance on safe sleep practices and appropriate infant positioning. Routine home cardiorespiratory monitoring is not generally recommended for low-risk BRUE. Nurses play an important role throughout management. Their responsibilities include monitoring respiratory patterns and vital signs, documenting the characteristics of each episode, implementing prescribed interventions, and evaluating treatment responses. They also support parents by explaining the monitoring process, demonstrating appropriate infant care, and addressing concerns before discharge. Effective management therefore requires early stabilization, identification of the underlying cause, appropriate treatment, continuous assessment, and family education.[16,22,23]

Differential Diagnosis

The differential diagnosis of apnea in infants is broad because several respiratory, cardiovascular, neurological, infectious, metabolic, and gastrointestinal conditions can produce episodes of interrupted breathing. Accurate identification of the underlying cause requires consideration of the infant's gestational age, medical history, clinical presentation, associated symptoms, and findings from physical examination and diagnostic investigations. Conditions that should be considered include aspiration syndrome, anemia of prematurity, botulism, bronchiolitis, bronchopulmonary dysplasia,

pediatric sleep apnea, croup, congenital heart disease with pulmonary hypertension, congestive heart failure, inborn errors of metabolism, influenza, laryngomalacia, factitious disorder imposed on another, and sepsis caused by serious bacterial or viral infections.[12,22] Respiratory disorders are important considerations. Bronchiolitis and influenza may cause apnea, particularly in young infants, and can present with cough, respiratory distress, nasal congestion, feeding difficulties, and oxygen desaturation. Croup may produce upper airway obstruction associated with a barking cough, stridor, and increased respiratory effort. Laryngomalacia can cause dynamic upper airway obstruction, particularly during feeding, crying, or lying supine. Pediatric sleep apnea should also be considered when episodes occur during sleep and are associated with snoring, abnormal respiratory patterns, or recurrent oxygen desaturation. Aspiration syndrome may produce respiratory compromise when milk, gastric contents, or other substances enter the airway. Premature infants require consideration of anemia of prematurity and bronchopulmonary dysplasia. Reduced oxygen-carrying capacity associated with anemia may contribute to cardiorespiratory instability, while bronchopulmonary dysplasia can cause persistent respiratory impairment and increased oxygen requirements. Neurological and neuromuscular conditions, including infant botulism, may also present with apnea. Botulism can produce progressive weaknesses, poor feeding, hypotonia, constipation, and impaired respiratory muscle function.

Cardiovascular disorders, particularly congenital heart disease associated with pulmonary hypertension and congestive heart failure, may cause hypoxemia, respiratory distress, poor feeding, sweating during feeding, and episodes of altered breathing. Inborn errors of metabolism should be considered when apnea occurs alongside lethargy, vomiting, poor feeding, seizures, or unexplained metabolic abnormalities. Sepsis caused by serious bacterial or viral infections is another critical diagnosis, especially when apnea is accompanied by temperature instability, lethargy, poor perfusion, or feeding intolerance. Factitious disorder imposed on another, formerly known as Munchausen syndrome by proxy, should be considered when episodes are unexplained, recurrent, or associated with inconsistent clinical observations. Assessment requires a careful, objective approach and appropriate safeguarding procedures. Distinguishing among these conditions is essential because their management strategies differ substantially. Early recognition of serious infectious, respiratory, cardiac, and metabolic causes helps prevent delayed treatment and associated complications. [12,22]

Consultations

The management of infant apnea may require consultation with several pediatric specialists, depending on the suspected underlying cause, clinical presentation, and response to initial treatment. A multidisciplinary approach helps identify conditions affecting the respiratory, neurological, cardiovascular, and upper airway systems. Pediatric otolaryngology consultation is appropriate when upper airway abnormalities are suspected. Conditions such as laryngomalacia, anatomical airway obstruction, and structural abnormalities may contribute to recurrent apnea, particularly during feeding or sleep. The otolaryngologist can assess airway anatomy and determine whether additional investigations or interventions are required. Pediatric neurology consultation may be necessary when apnea is associated with abnormal movements, altered responsiveness, hypotonia, seizures, or suspected neurological impairment. Neurological assessment can help identify central nervous system disorders affecting respiratory control. Pediatric cardiology consultation is indicated when episodes occur alongside bradycardia, cyanosis, abnormal cardiac findings, suspected arrhythmias, congenital heart disease, or pulmonary hypertension. Cardiac investigations may be required to evaluate structural and functional abnormalities. Pediatric pulmonology consultation is useful for infants with recurrent respiratory symptoms, persistent oxygen requirements, suspected chronic lung disease, or unexplained respiratory events. Pulmonary specialists can assess respiratory function and recommend appropriate respiratory support. Feeding-related apnea may require consultation with pediatric speech-language therapists, who can evaluate swallowing coordination and identify possible aspiration risks. Pediatric occupational therapists may also contribute by assessing feeding skills, positioning, sensory responses, and functional difficulties that interfere with safe feeding. The decision to involve specialists should be based on clinical findings rather than routine referral of every infant with apnea. The primary healthcare team should coordinate consultations, communicate relevant findings, and integrate specialist recommendations into a unified management plan. This approach supports timely diagnosis, reduces unnecessary investigations, and improves continuity of care during hospitalization and after discharge.

Patient Education

Parent and caregiver education is an essential component of managing infant apnea and reducing preventable risks. Families may experience considerable anxiety after witnessing an episode of interrupted breathing, changes in skin color, choking, or reduced responsiveness. Healthcare professionals should provide clear explanations of the infant's condition, the investigations performed, and the treatment plan. Parental reassurance should be based on the clinical assessment and should address individual concerns without minimizing the importance of recurrent or severe symptoms. Safe

sleep education is particularly important. Parents should be advised to place infants on their backs for every sleep period, use a firm and flat sleep surface, and keep the sleeping area free from pillows, loose blankets, toys, and other soft objects. Room-sharing without bed-sharing is recommended. Caregivers should also avoid exposing infants to tobacco smoke and other respiratory irritants. These practices support infant safety and reduce exposure to preventable sleep-related hazards. Parents and caregivers should learn to recognize concerning clinical signs, including persistent breathing difficulty, blue or pale skin, unusual lethargy, poor feeding, repeated choking, abnormal movements, and episodes of reduced responsiveness. They should understand when to seek urgent medical assessment and when emergency services are required. Recurrent apnea, prolonged breathing pauses, or episodes associated with poor responsiveness require prompt evaluation. Basic emergency response education can improve caregiver preparedness. Healthcare professionals should provide practical instruction on age-appropriate cardiopulmonary resuscitation and management of choking in infants. Training should include supervised demonstrations and opportunities for caregivers to practice the recommended techniques. Parents should also receive clear discharge instructions, relevant contact information, and guidance regarding follow-up appointments. Effective education helps families recognize emergencies, respond appropriately, and participate in the ongoing care of their infants.

Other Issues

Apneic episodes in infants may present as sudden events that cause considerable concern among parents and caregivers. Historically, some of these episodes were described as apparent life-threatening events (ALTEs). This term referred to an acute event that appeared frightening to the observer and involved one or more clinical features, including apnea, bradycardia, changes in skin color, choking, or gagging. Color changes could include cyanosis, pallor, and, less commonly, redness or plethora. These events may occur during sleep, feeding, or periods of wakefulness and require careful clinical assessment to determine their significance. In 2016, the American Academy of Pediatrics recommended discontinuing the use of ALTE as a diagnostic term and introduced brief resolved unexplained event (BRUE) as a more specific clinical classification. A BRUE describes a sudden, brief, and resolved episode in an infant younger than one year that involves at least one of the following features: cyanosis or pallor, absent or decreased breathing, marked change in muscle tone, or altered responsiveness. The event must remain unexplained after an appropriate history and physical examination. An important distinction is that not every episode of apnea qualifies as a BRUE. When a clear explanation is identified, such as choking during feeding, infection, or an underlying respiratory disorder, the event should be evaluated and managed according to that diagnosis. BRUE classification also requires consideration of risk factors. Infants who meet low-risk criteria may not require extensive investigations or hospital admission, whereas higher-risk presentations require individualized assessment and monitoring. Healthcare professionals should avoid using BRUE as a substitute for a comprehensive diagnostic evaluation. The classification does not identify the underlying mechanism or establish a specific disease. Instead, it provides a framework for describing unexplained events and guiding clinical decision-making. Accurate documentation of the event's duration, associated symptoms, circumstances, and recovery is essential. Clear communication with caregivers also helps prevent confusion between apnea, choking, normal variations in infant breathing, and clinically significant unexplained events.

Enhancing Healthcare Team Outcomes

Infant apnea has multiple possible causes and may occur intermittently, making early recognition and coordinated management essential. Recurrent apnea can result in intermittent hypoxia, particularly when episodes are prolonged or associated with bradycardia and oxygen desaturation. Persistent hypoxic episodes may contribute to cerebral injury and adversely affect neurodevelopmental outcomes.[13] Apnea associated with sepsis or neurological injury carries a high risk of mortality and requires prompt diagnosis and treatment.[23,24,25] If significant apnea remains untreated, affected infants may experience complications such as poor growth, impaired neurodevelopment, and cor pulmonale. Apnea is more common among infants with low birth weight and has been reported in approximately 1% to 7.5% of hospitalized infants. Improving healthcare team outcomes requires effective communication and clearly defined professional responsibilities. Neonatologists generally lead the diagnostic evaluation, establish treatment priorities, coordinate specialist involvement, and determine readiness for discharge.[26] Nurses provide continuous clinical observation, assess respiratory patterns, monitor oxygen saturation and heart rate, document apneic episodes, and implement prescribed interventions. They also provide tactile stimulation when indicated and communicate changes in the infant's condition to the medical team. Respiratory therapists contribute by assessing respiratory support requirements, managing oxygen delivery systems, and assisting with interventions such as CPAP.[14] Specialist involvement depends on the suspected cause. Neurologists may evaluate neurological abnormalities, cardiologists assess possible cardiac conditions, and gastroenterology or feeding

specialists may investigate gastrointestinal and swallowing-related problems.[14] Collaboration among these professionals supports accurate diagnosis and reduces delays in treatment. Discharge planning represents another important component of coordinated care. Before discharge, the team should establish appropriate observation criteria, confirm clinical stability, and determine whether medication or respiratory support remains necessary. Decisions about discontinuing caffeine or other treatment should follow the infant's clinical progress and applicable institutional protocols. Families should receive instructions on safe sleep, recognition of recurrent symptoms, emergency response, and scheduled follow-up.[13] Regular communication, accurate documentation, standardized monitoring protocols, and family participation can improve continuity of care. By combining medical expertise, nursing surveillance, respiratory support, specialist assessment, and caregiver education, the healthcare team can reduce preventable complications and support better outcomes for infants affected by apnea.

Conclusion:

Infant apnea is a clinical condition with diverse causes that require careful assessment and timely intervention. Its management depends on identifying the underlying etiology, recognizing associated clinical manifestations, and providing appropriate treatment. Premature infants may require caffeine therapy, continuous positive airway pressure, and cardiorespiratory monitoring, while term infants need investigations and interventions guided by their clinical condition. Early recognition of respiratory, neurological, cardiovascular, infectious, and metabolic causes helps prevent serious complications, including hypoxic injury and impaired neurodevelopment. Nurses play an essential role in monitoring respiratory patterns, documenting episodes, implementing prescribed interventions, and evaluating treatment responses. Collaboration among neonatologists, nurses, respiratory therapists, and relevant specialists supports coordinated care. Parental education regarding safe sleep, warning signs, emergency response, and follow-up is also necessary. Integrating clinical assessment, evidence-based treatment, continuous monitoring, and family participation can improve safety and outcomes for infants affected by apnea.

References:

1. Huang YS, Hsu JF, Paiva T, Chin WC, Chen IC, Guilleminault C. Sleep-disordered breathing, craniofacial development, and neurodevelopment in premature infants: a 2-year follow-up study. Sleep Med. 2019 Aug;60:20-25.
2. Kelly CE, Ooi WL, Yang JY, Chen J, Adamson C, Lee KJ, Cheong JLY, Anderson PJ, Doyle LW, Thompson DK. Caffeine for apnea of prematurity and brain development at 11 years of age. Ann Clin Transl Neurol. 2018 Sep;5(9):1112-1127.
3. Miura Y, Watanabe T, Uchida T, Nawa T, Endo N, Fukuzawa T, Ohkubo R, Takeyama J, Sasaki A, Hayasaka K. A novel PHOX2B gene mutation in an extremely low birth weight infant with congenital central hypoventilation syndrome and variant Hirschsprung's disease. Eur J Med Genet. 2019 Sep;62(9):103541.
4. Ramos-Fernández JM, Sánchez-Pérez M, Sánchez-González JM, Calvo-Cillán A, Moreno-Pérez D. Apnea in infants hospitalized with pertussis: Incidence and gestational smoking association. Pediatr Int. 2018 Oct;60(10):943-947
5. McFarlin A. What to Do when Babies Turn Blue: Beyond the Basic Brief Resolved Unexplained Event. Emerg Med Clin North Am. 2018 May;36(2):335-347.
6. Quitadamo P, Giorgio V, Zenzeri L, Baldassarre M, Cresi F, Borrelli O, Salvatore S. Apnea in preterm neonates: what's the role of gastroesophageal reflux? A systematic review. Dig Liver Dis. 2020 Jul;52(7):723-729.
7. Sanford JN, Lam DJ. Management of Obstructive Sleep Apnea in the Infant and Newborn. Otolaryngol Clin North Am. 2024 Jun;57(3):395-405.
8. Henderson-Smart DJ. The effect of gestational age on the incidence and duration of recurrent apnoea in newborn babies. Aust Paediatr J. 1981 Dec;17(4):273-6.
9. Regenbogen E, Zhang S, Yang J, Shroyer A, Zhu C, DeCristofaro J. Epidemiological trends among preterm infants with apnea. A twelve-year database review. Int J Pediatr Otorhinolaryngol. 2018 Apr;107:86-92.
10. Trachsel D, Erb TO, Hammer J, von Ungern-Sternberg BS. Developmental respiratory physiology. Paediatr Anaesth. 2022 Feb;32(2):108-117.
11. Erickson G, Dobson NR, Hunt CE. Immature control of breathing and apnea of prematurity: the known and unknown. J Perinatol. 2021 Sep;41(9):2111-2123.
12. Ginsburg D, Maken K, Deming D, Welch M, Fargo R, Kaur P, Terry M, Tinsley L, Ischander M. Etiologies of apnea of infancy. Pediatr Pulmonol. 2020 Jun;55(6):1495-1502.
13. Goodstein MH, Stewart DL, Keels EL, Moon RY., COMMITTEE ON FETUS AND NEWBORN, TASK FORCE ON SUDDEN INFANT DEATH SYNDROME. Transition to a Safe Home Sleep Environment for the NICU Patient. Pediatrics. 2021 Jul;148(1)
14. Sale SM. Neonatal apnoea. Best Pract Res Clin Anaesthesiol. 2010 Sep;24(3):323-36.
15. Zhao J, Gonzalez F, Mu D. Apnea of prematurity: from cause to treatment. Eur J Pediatr. 2011 Sep;170(9):1097-105.

16. Patrinos ME, Martin RJ. Apnea in the term infant. Semin Fetal Neonatal Med. 2017 Aug;22(4):240-244.
17. Pollack R, Lamichhane A. Term Infant with Apnea. Neoreviews. 2023 Oct 01;24(10):e670-e673.
18. Burstein B, Waterfield T, Umana E, Xie J, Kuppermann N. Prediction of Bacteremia and Bacterial Meningitis Among Febrile Infants Aged 28 Days or Younger. JAMA. 2026 Feb 03;335(5):425-433.
19. Moore BM, Roston TM, Laksman Z, Krahn AD. Updates on inherited arrhythmia syndromes (Brugada syndrome, long QT syndrome, CPVT, ARVC). Prog Cardiovasc Dis. 2025 Jul-Aug;91:130-143.
20. Choi HJ, Kim YH. Apparent life-threatening event in infancy. Korean J Pediatr. 2016 Sep;59(9):347-354.
21. Sen K, Agarwal R. More than meets the eye: infant presenting with hypoxic ischaemic encephalopathy. BMJ Case Rep. 2018 Apr 05;2018
22. Piumelli R, Davanzo R, Nassi N, Salvatore S, Arzilli C, Peruzzi M, Agosti M, Palmieri A, Paglietti MG, Nosetti L, Pomo R, De Luca F, Rimini A, De Masi S, Costabel S, Cavarretta V, Cremante A, Cardinale F, Cutrera R. Apparent Life-Threatening Events (ALTE): Italian guidelines. Ital J Pediatr. 2017 Dec 12;43(1):111.
23. Arane K, Claudius I, Goldman RD. Brief resolved unexplained event: New diagnosis in infants. Can Fam Physician. 2017 Jan;63(1):39-41.
24. Vliegenthart R, Miedema M, Hutten GJ, van Kaam AH, Onland W. High versus standard dose caffeine for apnoea: a systematic review. Arch Dis Child Fetal Neonatal Ed. 2018 Nov;103(6):F523-F529.
25. Hong HS, Lee JY. Intracranial hemorrhage in term neonates. Childs Nerv Syst. 2018 Jun;34(6):1135-1143.

Eichenwald EC., Committee on Fetus and Newborn, American Academy of Pediatrics. Apnea of Prematurity. Pediatrics. 2016 Jan;137(1).